

Analysis report No. 221027_AO2

Sample: 5-30 % CBD/ CBG tincture in HSO/OLI/SAL Oils

Sample type: oil tincture

Method: GC-FID

Sampling protocol: c.a. 130 - 250 mg sample in 5-15 ml acetone (p.a.) stirred for 20 minutes.

sample name (lot n°)	sample n°	CBD	Δ^9 -THC*	CBC	CBG	Δ^8 -THC	CBDv	THCv	CBN	CBT	CBL
		(wt.%)									
Natural 5% (CBGXE01-SAL4-05A)	221024.12	2.05	0.09	0.13	5.64	0.03	0.01	< 0.01	< 0.01	< 0.01	< 0.01
Natural 5% (APXE01-SAL4-05A)	221020.08	5.76	0.10	0.12	0.07	0.03	0.02	< 0.01	< 0.01	< 0.01	< 0.01
Chicken 5% (CBGXE01-HSO4-05A-HUH)	221020.07	2.37	0.10	0.15	5.37	0.03	0.01	< 0.01	< 0.01	< 0.01	< 0.01
Chicken 5% (APXE01-HSO4-05F-HUH)	221018.20	5.79	0.10	0.12	0.06	0.03	0.02	< 0.01	< 0.01	< 0.01	< 0.01
Beef 5% (CBGXE01-HSO4-05A-RIN)	221020.06	2.30	0.09	0.14	5.28	0.03	0.01	< 0.01	< 0.01	< 0.01	< 0.01
Beef 5% (APXE01-HSO4-05F-RIN)	221021.21	5.43	0.09	0.11	0.05	0.03	0.02	< 0.01	< 0.01	< 0.01	< 0.01

*Regulated cannabinoid (legal limit < 1 wt.% according to ordinance 812.121.11 of Swiss Federal Department of Home Affairs)

N.B.: Indicated values account for total amount of cannabinoids after complete decarboxylation of the material.



Bereket Tesfazghi
Chemist



Dr Jebril HADI
Head of Laboratory